

# The neural basis of functional brain imaging signals

David Attwell and Costantino Iadecola

The haemodynamic responses to neural activity that underlie the blood-oxygen-level-dependent (BOLD) signal used in functional magnetic resonance imaging (fMRI) of the brain are often assumed to be driven by energy use, particularly in presynaptic terminals or glia. However, recent work has suggested that most brain energy is used to power postsynaptic currents and action potentials rather than presynaptic or glial activity and, furthermore, that haemodynamic responses are driven by neurotransmitter-related signalling and not directly by the local energy needs of the brain. A firm understanding of the BOLD response will require investigation to be focussed on the neural signalling mechanisms controlling blood flow rather than on the locus of energy use.

The blood-oxygen-level-dependent (BOLD) magnetic resonance signal used in functional imaging of the brain reflects the loss of oxygen from haemoglobin, causing its iron to become paramagnetic, which influences the magnetic field experienced by protons in surrounding water molecules [1]. During neuronal activity, an increase of oxygen usage is followed within a few seconds by a larger fractional increase in blood flow and an increase in blood volume, resulting in a net decrease of the amount of deoxygenated haemoglobin present [2,3]. It is this that is detected in most BOLD imaging experiments. Positron-emission tomography (PET) imaging can also detect activity-induced increases in blood flow. Understanding the increased blood flow associated with neural activity, a relationship first proposed 100 years ago [4–8], is thus fundamental to understanding functional imaging. It is often assumed that this increase of blood flow is a direct consequence of the oxygen consumption or energy usage of the tissue [9–15] – an idea that we assess in this article. We initially consider which cellular processes the BOLD signal would reflect if blood flow were controlled by energy use, and then review whether blood flow changes and BOLD signals are in fact driven directly by energy consumption. We focus on blood flow changes because, although changes in the blood volume in activated brain areas affect the BOLD signal [16], the volume changes are probably a passive consequence of the blood-flow changes [9].

## Where is energy used in the brain?

If energy usage did control blood flow, then BOLD signals would reflect the activity of the cellular

processes consuming most energy. Brain energy usage has been attributed mainly to activity in presynaptic terminals [15] or to the energy needed to take up the neurotransmitter glutamate and convert it to glutamine in astrocytes [17]. These ideas have been investigated by constructing an energy budget based on the measured properties of individual ion channels and synapses [18]. This analysis concluded that most of the energy used by grey matter on signalling is, in fact, expended on reversing the ion fluxes underlying excitatory postsynaptic currents and action potentials (Fig. 1). In rodents, 47% of the energy used on signalling was predicted to support action potentials and 34% to support postsynaptic responses (>33% for postsynaptic currents and <1% for metabotropic responses). It was estimated that 10% supports the resting potential of neurons, with only 5% being used in glia (3% for the resting potential and 2% for glutamate recycling) and 3% used in synaptic terminals (<3% for  $\text{Ca}^{2+}$  fluxes and ~0.5% for vesicle filling and recycling). In primates, the greater number of synapses per neuron results in postsynaptic responses being predicted to be the dominant energy consumer (accounting for 74% of the energy used on signalling) [18], consistent with the high proportion (62%) of mitochondria found in dendrites [19]. Action potentials were predicted to consume 10% of the signalling energy used in primates, the neuronal resting potential to use 2%, glia to use 6% (5% for recycling glutamate and 1% for the resting potential) and presynaptic terminals to use 7%. Thus, assuming that experiments verify these predictions, if the BOLD signal from primates did directly reflect energy consumption it would be dominated by excitatory postsynaptic currents. In the following section, we argue that BOLD signals are, indeed, coupled to postsynaptic events – but not via energy use.

## Controversies over the locus of brain energy use

### *Energy used on action potentials*

The 10–47% of signalling energy predicted to be expended on action potentials [18] contrasts with a classic estimate by Creutzfeldt [20]. This estimate was based on heat generation by action potentials in peripheral nerves and suggested that only 0.3–3.0% of energy used in the brain was needed to support action potentials in human and cat cortex. However, Creutzfeldt's calculations employed a value of nerve heat production that was 6.6-fold lower than later measurements [21], he ignored the glial contribution to the nerve mass and, for cat cortex, he omitted a factor of ten for the action potential frequency. Redoing his calculation for human cortex, by dividing the heat that action potentials produce at 10 Hz ( $93 \mu\text{cal.g}^{-1}$  per action potential [21], assuming the same neural mass fraction as for peripheral nerve) by the total energy usage of grey matter ( $6870 \mu\text{cal.g}^{-1}\text{s}^{-1}$ , from a primate cortical glucose usage of  $57 \mu\text{mol per } 100 \text{ g.min}^{-1}$  [10]), suggests that

David Attwell\*  
Dept of Physiology,  
University College  
London, Gower Street,  
London, UK WC1E 6BT.  
\*e-mail: d.attwell@  
ucl.ac.uk

Costantino Iadecola  
Division of Neurobiology,  
Weill Medical College,  
Cornell University,  
411 East 69th Street,  
New York, NY 10021, USA.




Fig. 1. Schematic diagram of an excitatory neuron and a glial cell. The percentage of grey matter signalling energy predicted to be expended on different subcellular processes is shown for neurons (yellow) and glia (mauve) [18]. Predictions are given for rodents (first value) and primates (second value, calculated from rodent figures [18] by assuming a tenfold increase in synapse density per neuron with no other parameter changes).  $\text{Na}^+/\text{K}^+$  pump symbols illustrate the fact that almost all signalling energy is expended on reversing  $\text{Na}^+$  and  $\text{K}^+$  fluxes across the cell membrane (or reversing  $\text{Ca}^{2+}$  fluxes with  $\text{Na}^+/\text{Ca}^{2+}$  exchanger, which is indirectly powered by the  $\text{Na}^+/\text{K}^+$  pump). Presynaptic boutons represent release sites both on axon collaterals near the soma and on parts of the axon far from the soma in the grey matter. Mitochondrion icons illustrate sites of ATP production by oxidative phosphorylation: mitochondria in presynaptic terminals can provide ATP to  $\text{Na}^+/\text{K}^+$  pumps in the axon. The neurotransmitter glutamate (Glu) is recycled through uptake into glia and conversion to glutamine (Gln).

action potentials use 14% of the energy in primate grey matter. This is in line with more-detailed calculations [18].

#### Energy used on presynaptic terminals

The prediction that only 3–7% of energy is used in presynaptic terminals [18] contrasts with the observation that 23% of mitochondria are in synaptic terminals [19], which might suggest that presynaptic processes consume significant amounts of energy [15]. However, synaptic terminals are only  $5\ \mu\text{m}$  apart along cortical axons [22], implying a diffusion time for small molecules of only 25 ms. Thus, mitochondria in synaptic terminals might support energy use by axons, by providing the ATP needed to reverse the ion influx that drives axonal action potentials, as well as supporting presynaptic terminal activity (Fig. 1).

#### Energy used by astrocytes

Sibson *et al.* observed that rodent cortical glucose consumption is equal to the rate of glutamine formation from glutamate taken up into astrocytes [17] (Fig. 1). In the context of a model in which all ATP production in glia is glycolytic and glia then release lactate to neurons so that these can generate ATP by oxidative phosphorylation [12,13], it was concluded that the power needed for

'glutamate neurotransmitter cycling...accounts for >80% of total glucose oxidation' [17]. This is a discrepancy with the prediction that only 2% of brain energy supports glutamate recycling in glia [18]. Sibson *et al.* [17] apparently considered only glycolytically derived ATP, overlooking both the 94% of ATP produced by oxidative phosphorylation in mitochondria and the fact that glia need energy for tasks other than glutamate recycling, such as maintaining the resting potential [23,24]. This resulted in a 15-fold over-estimate of the fraction of ATP expended on glial glutamate recycling. Nevertheless, as most brain signalling energy is used by neurons, for postsynaptic currents and the action potentials that evoke transmitter release, one might expect the signalling energy used in a particular area to correlate with the (much smaller amount of) energy expended in the same area on recycling transmitter [18,25].

#### Energy used by inhibition

Many of the processes in inhibitory and excitatory neurons (e.g. the ion fluxes generating resting and action potentials) are identical, but inhibitory neurons and synapses are present at only one tenth of the density of their excitatory counterparts in the cortex [22,26]. Furthermore, the electrochemical gradient down which  $\text{Cl}^-$  moves postsynaptically at inhibitory synapses is not as steep as that down which  $\text{Na}^+$  moves at excitatory synapses, implying that less energy expenditure is required to pump the ions back. Waldvogel *et al.* [27] reported that inhibition evokes no BOLD signal, and concluded that inhibitory synapses use less energy than excitatory synapses. However, in their experiments, inhibition was associated with the generation of fewer action potentials, so the change of energy use evoked by their stimuli does not reflect just inhibitory synaptic activity. In fact, increased glucose usage is associated with inhibition of hippocampal pyramidal and auditory cells [28,29], although it is not clear whether the energy is needed to drive inhibitory synaptic currents or to power the action potentials that generate inhibitory transmitter release.

#### What controls cerebral blood flow?

##### Energy use does not directly increase blood flow

The preceding discussion suggests that, in primates, most of the increased energy use generated by neuronal activity is expended on reversing the ion movements that generate excitatory postsynaptic currents, with a smaller fraction being devoted to reversing the ion movements that generate action potentials. Does this energy usage *directly* control the blood flow to active regions of the brain and, hence, determine the functional imaging signal from those areas? Recent evidence suggests not. Although changes in cerebral blood flow (CBF) correlate with oxygen usage during functional activation of the




Fig. 2. Schematic diagram of the wiring of the cerebellar cortex, with input excitatory axons (climbing fibre and mossy fibre inputs) shown in purple, inhibitory interneurons (Golgi, basket and stellate cells) in blue, excitatory granule interneurons in lime green, and the output Purkinje cell in dark turquoise. Specimen Purkinje cell action potentials are from Ref. [66]. Crossed arrow indicates that blood flow is not controlled by factors reflecting the spiking output of Purkinje cells [49,50]. Other arrows indicate that vasodilators released from stellate cells [53] or from the presynaptic terminals of the parallel fibres [52] probably control blood flow, so that blood-flow changes reflect information processing activity within the cerebellar cortex.

brain [9], this need not imply that oxygen or energy usage directly regulate the blood flow. Indeed, the available evidence suggests that oxygen and glucose delivery to the tissue is not maintained by oxygen or glucose lack increasing the CBF [30,31]. Neither can signals related to energy production, such as  $\text{CO}_2$  and  $\text{H}^+$ , account for the changes in CBF evoked by neural activity [32,33]. How, then, can CNS tissue maintain an energy supply adequate to meet its changing requirements? Interestingly, although the oxygen usage associated with neuronal activity must colocalize with the activity, the subsequent increase of blood flow occurs in a larger area [34], suggesting that blood flow is controlled by factors other than a lack of energy.

#### *Control of local blood flow by fast neurotransmitters*

In the cerebellum, hippocampus and neocortex, there is increasing evidence that blood flow is controlled locally by glutamate and, perhaps, by GABA. In the cerebellar cortex, activating the excitatory parallel or climbing fibre inputs leads to an increased blood flow that is blocked by inhibitors of non-NMDA glutamate receptors, nitric oxide synthase (NOS) and adenosine receptors [35–37]. In addition, exogenous glutamate mimics the vascular effects of parallel fibre stimulation [38]. Similarly, in neocortex and hippocampus, exogenous glutamate or NMDA dilates pial arterioles and/or precapillary microvessels – an effect attenuated by NOS

inhibitors [39–41]. The increase in CBF produced by activation of the somatosensory cortex is also attenuated by inhibition or genetic deletion of cyclooxygenase-2, an enzyme that synthesizes prostanoids from arachidonic acid and is associated with glutamatergic synapses [42,43]. These data suggest that a glutamate-evoked  $\text{Ca}^{2+}$  influx in postsynaptic neurons activates the production of NO, adenosine and arachidonic acid metabolites. These agents, in turn, produce a vasodilation reflecting both the activity of neurons presynaptic to the cells releasing NO, adenosine and prostanoids, and the level of depolarization of the postsynaptic cell (which will alter  $\text{Mg}^{2+}$ -block of NMDA receptors and the resulting  $\text{Ca}^{2+}$  influx). Interestingly, in the cerebellum the blood vessel dilation produced locally by parallel fibre activation somehow propagates to more distant blood vessels [44], providing a possible explanation for the observation that blood flow increases over a larger area than that in which there is neuronal activity [34].

Activity-evoked rises in extracellular  $\text{K}^+$  concentration ( $[\text{K}]_o$ ) have also been shown to increase blood flow [45,46], but they have a less of an effect than neural activity. In addition, much of the effect of the  $[\text{K}]_o$  increase could be via  $\text{K}^+$ -depolarizing neurons and, thus, releasing glutamate or evoking a  $\text{Ca}^{2+}$  influx via voltage-gated channels, which leads to release of nitric oxide (NO) [47].

In the light of the controversy over whether inhibition contributes to brain energy use or to functional imaging signals, it is interesting that exogenous GABA, acting via  $\text{GABA}_A$  receptors, dilates precapillary microvessels in hippocampus and neocortex, and that blockage of  $\text{GABA}_A$  receptors produces a constriction [48]. However, in the cerebellar cortex, GABA is not involved in activation-induced increases in CBF [36,49].

#### *Local blood flow and the BOLD signal do not correlate with principal neuron spike rate*

In the cerebellar cortex, it is not possible to relate neurally evoked blood flow changes (and, therefore, BOLD signals) to an increase in firing of the principal output neurons, the Purkinje cells. Indeed, parallel fibre stimulation increases blood flow but inhibits Purkinje cell spiking [49,50]. Similarly, in visual cortex the BOLD signal correlates somewhat better with synaptically evoked field potentials than it does with the spiking output of the area [51]. It is essential to determine whether these results, which could reflect NOS being located both in excitatory and inhibitory neurons [52], extend to other brain areas. In the cerebellar cortex of mice with cyclin D2 knocked out, which decreases the number of inhibitory NOS-containing stellate cells, there is a reduction in the blood flow response to neural activity, suggesting that stellate cells play a significant role in controlling cerebellar blood flow [53]. Although knockout of cyclin D2 has effects

other than deleting stellate cells (e.g. decreasing the number of granule cells), if activation of NOS in stellate cells was the main factor controlling cerebellar cortical blood flow, then blood flow or BOLD signals from the cerebellum might correlate best with the firing rate of parallel fibres (i.e. granule cells) or with the firing of the inhibitory stellate cells. The blood flow or BOLD signals would, thus, reflect information processing within the cerebellar cortex, rather than the firing of the output Purkinje cells or of the input mossy fibres (Fig. 2).

#### *Global control of blood flow by dedicated intrinsic neural networks*

In the cerebral cortex, dopamine, noradrenaline and serotonin are present in neurons innervating microvessels and produce vasoconstriction [54–57]. Furthermore, cholinergic axons from the basal forebrain have close associations with neocortical arterioles and capillaries, and can produce vasodilation [58,59]. Thus, in addition to the spatially restricted control of blood flow provided by fast transmitters, there might be a more widespread regulation by dopaminergic fibres from the mesencephalic ventral tegmental area, noradrenergic fibres from the locus coeruleus, serotonergic fibres from the raphe nucleus, and cholinergic fibres from the basal forebrain. These projections are diffuse, and could increase the blood flow to a wide cortical area when a particular type of task is being carried out. Furthermore, activation of some neural pathways, such as those passing through the cerebellar fastigial nucleus or originating in area C1 of the rostral ventrolateral medulla, leads to a general rise in CBF without a change of metabolism [60]. This is mediated, in part, by muscarinic ACh receptors and release of NO [60,61], and raises the possibility of a global change of CBF independent of local energy needs.

#### **Functional imaging of neuropsychiatric conditions**

The findings that amines could be involved in CBF control and that CBF can be dissociated from energy utilization have important implications for the interpretation of functional magnetic resonance imaging (fMRI) data in disease states. Functional imaging is increasingly being used to investigate

brain function in conditions with altered amine function, such as schizophrenia, Parkinson's disease, attention-deficit hyperactivity disorder or infusion of drugs affecting amine receptors or transporters [62–65]. However if dopamine, noradrenaline and serotonin modulate the blood flow changes that generate the BOLD signal, then in such conditions it might be hard to determine whether any observed alteration of the BOLD signal is due to an effect of altered amine function on cortical processing, or simply an alteration of the neural control of blood flow. In such work, control experiments will need to be devised to test how the amine disorder affects neural control of CBF, independently of affecting neural processing.

#### **Conclusions**

The available evidence indicates that the haemodynamic response to neural activity is not initiated by signals arising from the energy deficit of the tissue but, rather, is driven locally by fast glutamate-mediated signalling processes, and more globally by amine- and ACh-mediated neural systems. Accordingly, the BOLD effect used in functional brain imaging should be interpreted as a reflection of neuronal signalling and not as a locus of increased energy utilization. BOLD signals could, as in cerebellum, reflect the neural processing occurring within a brain area, rather than the output from or input to that area. Consequently, a change of processing with no net change of energy usage could lead to a BOLD signal – and a change of spiking output with no change to the signalling systems controlling blood flow could fail to generate a BOLD signal. Although sites of increased neural activity can often colocalize with areas of increased metabolism, CBF and energy utilization can be dissociated and should be considered as the results of processes operating in parallel. The view that the haemodynamic response is coupled to signalling processes represents a conceptual shift from the traditional idea that the energy demands of the tissue directly determine the flow increase associated with neural activation. In summary, we suggest that understanding the BOLD response is a signalling problem, not an energy problem.

#### **Acknowledgements**

We thank Simon Laughlin for extended discussions on these issues, and Tony David and Chris Frith for comments on the manuscript. Supported by the Wellcome Trust and a Wolfson-Royal Society Award (DA) and the NIH (CI).

#### **References**

- Ogawa, S. *et al.* (1990) Brain magnetic resonance imaging with contrast dependent on blood oxygenation. *Proc. Natl. Acad. Sci. U. S. A.* **87**, 9868–9872
- Fox, P.T. and Raichle, M.E. (1986) Focal physiological uncoupling of cerebral blood flow and oxidative metabolism during somatosensory stimulation in human subjects. *Proc. Natl. Acad. Sci. U. S. A.* **83**, 1140–1144
- Malonek, D. *et al.* (1997) Vascular imprints of neuronal activity: relationships between the dynamics of cortical blood flow, oxygenation, and volume changes following sensory stimulation. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 14826–14831
- Mosso, A. (1881) *Ueber den Kreislauf des Blutes im menschlichen Gehirn*, von Veit, Leipzig
- Roy, C. and Sherrington, C. (1890) On the regulation of the blood supply of the brain. *J. Physiol.* **11**, 85–100
- Fulton, J.F. (1928) Observations upon the vascularity of the human occipital lobe during visual activity. *Brain* **LI**, 310–320
- Schmidt, C. and Hendrix, J. (1938) Action of chemical substances on cerebral blood vessels. *Res. Publ. Assoc. Res. Nerv. Ment. Dis.* **18**, 229–276
- Raichle, M.E. (1998) Behind the scenes of functional brain imaging: a historical and physiological perspective. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 765–772
- Hoge, R.D. *et al.* (1999) Linear coupling between cerebral blood flow and oxygen consumption in activated human cortex. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 9403–9408
- Clark, J.B. and Sokoloff, L. (1999) Circulation and energy metabolism of the brain. In *Basic Neurochemistry* (6th edn; Siegel, G.J. *et al.* eds), pp 637–669, Lippincott–Raven
- Shulman, R.G. and Rothman, D.L. (1998) Interpreting functional imaging studies in terms of neurotransmitter cycling. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 11993–11998

- 12 Magistretti, P.J. *et al.* (1999) Energy on demand. *Science* 283, 496–497
- 13 Magistretti, P.J. and Pellerin, L. (1999) Cellular mechanisms of brain energy metabolism and their relevance to functional brain imaging. *Philos. Trans. R. Soc. London Ser. B* 354, 1155–1163
- 14 Shin, C. (2000) Neurophysiologic basis of functional neuroimaging: animal studies. *J. Clin. Neurophysiol.* 17, 2–9
- 15 Jueptner, M. and Weiller, C. (1995) Does measurement of regional cerebral blood flow reflect synaptic activity? – Implications for PET and fMRI. *Neuroimage* 2, 148–156
- 16 Mandeville, J.B. *et al.* (1999) MRI measurement of the temporal evolution of relative CMRO<sub>2</sub> during rat forepaw stimulation. *Magn. Reson. Med.* 42, 944–951
- 17 Sibson, N.R. *et al.* (1998) Stoichiometric coupling of brain glucose metabolism and glutamatergic neuronal activity. *Proc. Natl. Acad. Sci. U. S. A.* 95, 316–321
- 18 Attwell, D. and Laughlin, S.B. (2001) An energy budget for signalling in the grey matter of the brain. *J. Cereb. Blood Flow Metab.* 21, 1133–1145
- 19 Wong-Riley, M.T.T. (1989) Cytochrome oxidase: an endogenous metabolic marker for neuronal activity. *Trends Neurosci.* 12, 94–101
- 20 Creutzfeldt, O.D. (1975) Neurophysiological correlates of different functional states of the brain. In *Alfred Benzon Symposium VIII* (Ingvar D.H. and Lassen N.A., eds), pp 21–46, Academic Press
- 21 Ritchie, J.M. (1973) Energetic aspects of nerve conduction: the relationships between heat production, electrical activity and metabolism. *Prog. Biophys. Mol. Biol.* 26, 149–187
- 22 Braitenberg, V. and Schüz, A. (1998) *Cortex: Statistics and Geometry of Neuronal Connectivity* (2nd edn), Springer
- 23 Hertz, L. *et al.* (1999) Energy for neurotransmission. *Science* 285, 639a
- 24 Chih, C.P. *et al.* (2001) Do neurons really use lactate rather than glucose? *Trends Neurosci.* 24, 573–578
- 25 Bonvento, G. *et al.* (2002) Does glutamate image your thoughts? *Trends Neurosci.* 25, 359–364
- 26 Abeles, M. (1991) *Corticonics: Neural Circuits of the Cerebral Cortex*, Cambridge University Press
- 27 Waldvogel, D. *et al.* (2000) The relative metabolic demand of excitation and inhibition. *Nature* 406, 995–998
- 28 Ackermann, R.F. (1984) Increased glucose metabolism during long-duration recurrent inhibition of hippocampal pyramidal cells. *J. Neurosci.* 4, 251–264
- 29 Nudo, R.J. and Masterton, R.B. (1986) Stimulation-induced [<sup>14</sup>C]2-deoxyglucose labeling of synaptic activity in the central auditory system. *J. Comp. Neurol.* 245, 553–565
- 30 Mintun, M.A. *et al.* (2001) Blood flow and oxygen delivery to human brain during functional activity: theoretical modeling and experimental data. *Proc. Natl. Acad. Sci. U. S. A.* 98, 6859–6864
- 31 Powers, W.J. *et al.* (1996) Effect of stepped hypoglycemia on regional cerebral blood flow response to physiological brain activation. *Am. J. Physiol.* 270, H554–H559
- 32 Astrup, J. *et al.* (1978) Evidence against H<sup>+</sup> and K<sup>+</sup> as main factors for the control of cerebral blood flow: a microelectrode study. In *Cerebral vascular smooth muscle and its control* (K. Elliott and M. O'Connor, eds), pp 313–337, Elsevier
- 33 Pinard, E. *et al.* (1984) Blood flow compensates oxygen demand in the vulnerable CA3 region of the hippocampus during kainate-induced seizures. *Neuroscience* 13, 1039–1049
- 34 Malonek, D. and Grinvald, A. (1996) Interactions between electrical activity and cortical microcirculation revealed by imaging spectroscopy: implications for functional brain mapping. *Science* 272, 551–554
- 35 Akgören, N. *et al.* (1994) Importance of nitric oxide for local increases of blood flow in rat cerebellar cortex during electrical stimulation. *Proc. Natl. Acad. Sci. U. S. A.* 91, 5903–5907
- 36 Li, J. and Iadecola, C. (1994) Nitric oxide and adenosine mediate vasodilation during functional activation in cerebellar cortex. *Neuropharmacology* 33, 1453–1461
- 37 Yang, G. *et al.* (1999) Nitric oxide is the predominant mediator of cerebellar hyperemia during somatosensory activation in rats. *Am. J. Physiol.* 277, R1760–R1770
- 38 Yang, G. and Iadecola, C. (1996) Glutamate microinjections in cerebellar cortex reproduce cerebrovascular effects of parallel fibre stimulation. *Am. J. Physiol.* 271, R1568–R1575
- 39 Faraci, F.M. and Breese, K.R. (1993) Nitric oxide mediates vasodilation in response to activation of N-methyl-D-aspartate receptors in brain. *Circ. Res.* 72, 476–480
- 40 Fergus, A. and Lee, K.S. (1997) Regulation of cerebral microvessels by glutamatergic mechanisms. *Brain Res.* 754, 35–45
- 41 Lovick, T. (1999) Neurovascular relationships in hippocampal slices: physiological and anatomical studies of mechanisms underlying flow-metabolism coupling in intraparenchymal microvessels. *Neuroscience* 92, 47–60
- 42 Niwa, K. *et al.* (2000) Cyclooxygenase-2 contributes to functional hyperemia in whisker-barrel cortex. *J. Neurosci.* 20, 763–770
- 43 Kaufmann, W.E. *et al.* (1996) COX-2, a synaptically induced enzyme, is expressed by excitatory neurons at postsynaptic sites in rat cerebral cortex. *Proc. Natl. Acad. Sci. U. S. A.* 93, 2317–2321
- 44 Iadecola, C. *et al.* (1997) Local and propagated vascular responses evoked by focal synaptic activity in cerebellar cortex. *J. Neurophysiol.* 78, 651–659
- 45 Iadecola, C. and Kraig, R.P. (1991) Focal elevations in neocortical interstitial K<sup>+</sup> produced by stimulation of the fastigial nucleus in rat. *Brain Res.* 563, 273–277
- 46 Caesar, K. *et al.* (1999) Modification of activity-dependent increases in cerebellar blood flow by extracellular potassium in anaesthetized rats. *J. Physiol.* 520, 281–292
- 47 Dreier, J.P. *et al.* (1995) Nitric oxide modulates the CBF response to increased extracellular potassium. *J. Cereb. Blood Flow Metab.* 15, 914–919
- 48 Fergus, A. and Lee, K.S. (1997) GABAergic regulation of cerebral microvascular tone in the rat. *J. Cereb. Blood Flow Metab.* 17, 992–1003
- 49 Mathiesen, C. *et al.* (1998) Modification of activity-dependent increases of cerebral blood flow by excitatory synaptic activity and spikes in rat cerebellar cortex. *J. Physiol.* 512, 555–566
- 50 Lauritzen, M. (2001) Relationship of spikes, synaptic activity and local changes of cerebral blood flow. *J. Cereb. Blood Flow Metab.* 21, 1367–1383
- 51 Logothetis, N.K. *et al.* (2001) Neurophysiological investigation of the basis of the fMRI signal. *Nature* 412, 150–157
- 52 Southam, E. *et al.* (1992) Sources and targets of nitric oxide in rat cerebellum. *Neurosci. Lett.* 137, 241–244
- 53 Yang, G. *et al.* (2000) Stellate neurons mediate functional hyperemia in the cerebellar molecular layer. *J. Neurosci.* 20, 6968–6973
- 54 Krimer, L.S. *et al.* (1998) Dopaminergic regulation of cerebral cortical microcirculation. *Nat. Neurosci.* 1, 286–289
- 55 Raichle, M.E. *et al.* (1975) Central noradrenergic regulation of cerebral blood flow and vascular permeability. *Proc. Natl. Acad. Sci. U. S. A.* 72, 3726–3730
- 56 Cohen, Z. *et al.* (1997) Astroglial and vascular interactions of noradrenaline terminals in the rat cerebral cortex. *J. Cereb. Blood Flow Metab.* 17, 894–904
- 57 Cohen, Z. *et al.* (1996) Serotonin in the regulation of brain microcirculation. *Prog. Neurobiol.* 50, 335–362
- 58 Chedotal, A. *et al.* (1994) Light and electron microscopic immunocytochemical analysis of the neurovascular relationships of choline acetyltransferase and vasoactive intestinal polypeptide nerve terminals in the rat cerebral cortex. *J. Comp. Neurol.* 343, 57–71
- 59 Sato, A. and Sato, Y. (1992) Regulation of regional cerebral blood flow by cholinergic fibers originating in the basal forebrain. *Neurosci. Res.* 14, 242–274
- 60 Reis, D.J. and Golanov, E.V. (1997) Autonomic and vasomotor regulation. *Int. Rev. Neurobiol.* 41, 121–149
- 61 Iadecola, C. (1992) Nitric oxide participates in the cerebrovasodilation elicited from cerebellar fastigial nucleus. *Am. J. Physiol.* 263, R1156–R1161
- 62 Sabatini, U. *et al.* (2000) Cortical motor reorganization in akinetic patients with Parkinson's disease: a functional MRI study. *Brain* 123, 394–403
- 63 Vaidya, C.J. *et al.* (1998) Selective effects of methylphenidate in attention deficit hyperactivity disorder: a functional magnetic resonance study. *Proc. Natl. Acad. Sci. U. S. A.* 95, 14494–14499
- 64 Nguyen, T.V. *et al.* (2000) Detection of the effects of dopamine receptor supersensitivity using pharmacological MRI and correlations with PET. *Synapse* 36, 57–65
- 65 Li, S.-J. *et al.* (2000) Cocaine administration decreases functional connectivity in human primary visual and motor cortex as detected by functional MRI. *Magn. Reson. Med.* 43, 45–51
- 66 Hamann, M. *et al.* (2002) Tonic and spillover inhibition of granule cells control information flow through cerebellar cortex. *Neuron* 33, 625–633